

COZEAN PELVIC DYSFUNCTION SCREENING PROTOCOL

INSTRUCTIONS: CHECK ALL THAT APPLY

- ☐ I sometimes have pelvic pain (in genitals, perineum, pubic or bladder area, or pain with urination) that exceeds a '3' on a 1-10 pain scale, with 10 being the worst pain imaginable
- ☐ I can remember falling onto my tailbone, lower back, or buttocks (even in childhood)
- ☐ I sometimes experience one or more of the following urinary symptoms
 - Accidental loss of urine
 - Feeling unable to completely empty my bladder
 - Having to void within a few minutes of a previous void
 - Pain or burning with urination
 - Difficulty starting or frequent stopping/starting of urine stream
- ☐ I often or occasionally have to get up to urinate two or more times at night
- ☐ I sometimes have a feeling of increased pelvic pressure or the sensation of my pelvic organs slipping down or falling out
- ☐ I have a history of pain in my low back, hip, groin, or tailbone or have have scatica
- ☐ I sometimes experience one or more of the following bowel symptoms
 - Loss of bowel control
 - Feeling unable to completely empty my bowels
 - Straining or pain with a bowel movement
 - Difficulty initiating a bowel movement
- ☐ I sometimes experience pain or discomfort with sexual activity or intercourse
- ☐ Sexual activity increases one or more of my other symptoms
- ☐ Prolonged sitting increases my symptoms

**IF YOU CHECKED 3 OR MORE CIRCLES,
PELVIC FLOOR DYSFUNCTION IS LIKELY. YOU MAY BENEFIT FROM AN
ASSESSMENT FROM A PELVIC FLOOR REHABILITATION (PT/OT) PROVIDER.**

© Nicole Cozean PT, DPT, WCS. All rights reserved.